

**CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM**  
**maintained by the**  
**CALIFORNIA DEPARTMENT OF FISH AND GAME**  
**and supported by the**  
**CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP**  
**Database Version 8.1 (2005)**

---

B154 Snowy Plover *Charadrius alexandrinus*  
Family: Charadriidae Order: Charadriiformes Class: Aves

Written by: M. Rigney  
Reviewed by: L. Mewaldt  
Edited by: S. Granholm, R. Duke  
Updated by: CWHR Program Staff, February 2005

#### DISTRIBUTION, ABUNDANCE, AND SEASONALITY

In fall and winter, common on sandy marine and estuarine shores, uncommon at salt ponds, and rare at the Salton Sea. Nests locally in these same habitats from April through August, but the major nesting habitat now appears to be on salt pond levees (Cogswell 1977). Inland nesting areas occur at the Salton Sea, Mono Lake, and at isolated sites on the shores of alkali lakes in northeastern California, in the Central Valley, and southeastern deserts (Jurek and Leach 1973, Page et al. 1979, 1983, Garrett and Dunn 1981).

#### SPECIFIC HABITAT REQUIREMENTS

**Feeding:** Gleans insects and amphipods from the dry sand of upper beaches along the coast (Cogswell 1977). Occasionally forages in wet sand for young sand crabs. At salt ponds and alkali lakes, feeds primarily on brine flies.

**Cover:** Crouches motionless on sandy substrate and relies on camouflage for cover.

**Reproduction:** Requires a sandy, gravelly or friable soil substrate for nesting. Nests are shallow depressions in the sand or soil, sometimes lined with small pebbles, glass fragments, or gravel. Frequently locates nest near or under objects such as driftwood, rocks, or defoliated bushes. Nests also may be on barren ground with no nearby cover (Bent 1929, Jurek and Leach 1973).

**Water:** May stand in water for cooling at some of the hotter inland nesting areas.

**Pattern:** Nests, feeds, and takes cover on sandy or gravelly beaches along the coast, on estuarine salt ponds, alkali lakes, and at the Salton Sea.

#### SPECIES LIFE HISTORY

**Activity Patterns:** Yearlong, diurnal activity.

**Seasonal Movements/Migration:** Beginning in July and August, may move from northwest Oregon to as far as Baja California. Remains on wintering grounds from September through March. Smaller numbers remain year-round at the Salton Sea and at salt ponds on San Francisco Bay (Cogswell 1977).

**Home Range:** During the breeding season, adult generally does not wander far from the nest (Page et al. 1977). At Mono Lake, breeding adult feeds at freshwater seeps up to 1.5 km (0.94 mi) away from the nest site (Page et al. 1995).

**Territory:** Nesting density and territorial defense appear to depend on predators. Nesting density was 1 nest/6 ha (15 ac) at Mono Lake, where predatory pressure was high, while density was 20 nests/6 ha (15 ac) at Monterey Bay, where predators were infrequent (Page et al. 1983).

**Reproduction:** Present at nesting sites from April through August. Solitary nester; clutch size 2-6, average = 3 eggs. Incubation, mostly by the male, lasts approximately 24 days. The young are precocial, following the adults to feeding areas within 1 day. The young are agile and able to avoid predators within 2 days after hatching. Fledging age reported as 29-47 days (Ehrlich et al. 1988).

**Niche:** Gulls, ravens, coyotes, and skunks are important predators of adults, eggs, and young at Mono Lake (Page et al. 1983). The historical nesting area for snowy plovers, sandy marine beaches, has brought them into constant contact with humans using these areas for recreation. The increase in nesting populations at salt ponds and other inland areas is opportunistic and compensates somewhat for increased human disturbance at the coastal nesting sites (Jurek and Leach 1973, Cogswell 1977).

## REFERENCES

- Bent, A. C. 1929. Life histories of North American shorebirds. Part 2. U.S. Natl. Mus. Bull. 146. 412pp.
- Cogswell, H. L. 1977. Water birds of California. Univ. California Press, Berkeley. 399pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The birder's handbook. Simon and Schuster, New York. 785pp.
- Garrett, K., and J. Dunn. 1981. Birds of southern California. Los Angeles Audubon Soc. 408pp.
- Jurek, R. M., and H. R. Leach. 1973. California shorebird study: Accelerated research program on shore and upland migratory game birds. Calif. Dept. Fish and Game, Sacramento. P. R. project W-54-R. Final Rep. 233pp.
- Page, G. W., J. S. Warriner, J. C. Warriner, and P. W. C. Paton. 1995. Snowy Plover (*Charadrius alexandrinus*). In The Birds of North America No. 154 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C.
- Page, G. W., D. F. Winkler, and C. W. Swarth. 1979. Breeding success of snowy plovers at Mono Lake, California. U. S. Fish and Wildl. Serv., Portland OR. Refuge and Wildl. Resour. Dep. Rep. 21pp.
- Page, G. W., L. E. Stenzel, D. W. Winkler, and C. W. Swarth. 1983. Spacing out at Mono Lake: breeding success, nest density, and predation in the snowy plover. Auk 100:13-24.
- Page, G. W., L. E. Stenzel, and C. M. Wolfe. 1979. Aspects of the occurrence of shorebirds on a central California estuary. Pages 15-32 in F. A. Pitelka, ed. Shorebirds in marine environments. Studies in Avian Biol. No. 2. Cooper Ornithol. Soc. Lawrence, KA. 261pp.
- Page, G. W., J. S. Warriner, J. C. Warriner, R. M. Halbresen, and S. C. Peaslee. 1977. Status of the snowy plover on the northern California coast. Calif. Dep. Fish and Game, Sacramento. Nongame Wildl. Invest. Rep. 25pp.
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. Calif. Dept. of Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1. 54pp.